



ELSEVIER

Cancer Genetics and Cytogenetics ■ (2005) ■

 CANCER GENETICS
 AND
 CYTOGENETICS

Letter to the editor

B lymphoid or myeloid lineage identity of cell lines derived from chronic myeloid leukemia blast crisis

Chronic myeloid leukemia (CML) develops as a triphasic disease, ultimately leading to fatal blast crisis unless successfully treated [1]. Recent data have identified DOK1/DOK2 [2,3] and β -catenin [4] as important targets in the transition of chronic phase CML into blast crisis, but the molecular mechanism determining blastoid transformation of CML cells is not fully understood. In almost all instances, CML cells carry an oncogenic *BCR-ABL1* gene rearrangement resulting from the t(9;22)(q34;q11) translocation [5]. Studies in murine and human hematopoietic progenitor cells have shown that expression of a *BCR-ABL1* fusion gene is required and sufficient to drive malignant transformation of myeloid progenitor cells [6,7]. Also, B cell precursors can be transformed by oncogenic BCR-ABL1 kinase activity [6]. Indeed, *BCR-ABL1* gene rearrangements represent the most frequent recurrent genetic aberrations in adult B lineage acute lymphoblastic leukemias and are detected in ~25% of the cases [5]. Involvement of the major breakpoint cluster region (MBCR) within the *BCR* gene leading to the expression of a fusion protein of 210 kD (P210 BCR-ABL1) is characteristic for CML (>99%) [5]. In ~70% of *BCR-ABL1*⁺ B cell lineage acute lymphoblastic leukemia, the leukemia cells carry *BCR-ABL1* fusion genes rearranged within the minor breakpoint cluster region (mBCR) encoding a smaller BCR-ABL1 fusion protein of 190 kD [5]. Recent work has shown that B lymphoid and myeloid cells transformed by P210 BCR-ABL1 markedly differ in their signaling requirements for sustained growth and survival [8]. The reason for different distribution of *BCR-ABL1* breakpoints in CML and B cell lineage acute lymphoblastic leukemia, however, is still currently unclear.

Of note, a significant fraction of cases of CML exhibit a B lymphoid phenotype upon progression into blast crisis. The phenotypic lineage conversion in these cases remains enigmatic. So-called *lymphoid blast crisis cells* typically express P210 BCR-ABL1 fusion proteins, suggesting that they represent the lymphoid outgrowth of a common initial CML clone. Alternatively, a B lymphoid clone unrelated to chronic phase CML may outcompete CML cells at a later stage of the disease and then emerge as lymphoid blast crisis. To date, it is not resolved whether lymphoid blast crisis cells share a common clonal origin with chronic phase CML or whether they represent the progeny of a B lymphoid

tumor clone independent from chronic phase CML. It is also unclear whether B cell lineage leukemia cells expressing p210 BCR-ABL1 can develop as a primary acute lymphoblastic leukemia or only secondary to CML in the context of lymphoid blast crisis.

In either case, however, it is important to distinguish between myeloid and B lymphoid cells in the development of CML blast crisis. For instance, homozygous deletions within the *CDKN2A* (alias *INK4A*) locus can be detected in ~50% of the cases of B lymphoid blast crisis, but not in myeloid blast crisis [9]. Moreover, B lymphoid but not myeloid cells transformed by P210 BCR-ABL1 critically depend on SRC-kinase activity [8].

In the meantime, a large panel of CML blast crisis-derived cell lines is available [10], including both myeloid and lymphoid variants of CML blast crisis. Among them two cell lines, BV173 and NALM1, are B lymphoid in origin [11–13] but often referred to as “CML cell lines” [14–29]. The recent study by Pelz and colleagues [30] comprehensively reanalyzed the karyotype of NALM1 cells to establish this cell line as a “CML control.” Although we do not question the accuracy of their cytogenetic analysis, we do disagree with their suggestion to use NALM1 cells as a control in studies related to CML cells. Unlike CML cells, NALM1 cells carry an in-frame immunoglobulin heavy chain V_H3.9–D_H2.21–J_H3 gene rearrangement on one allele of the *IGH* locus and a D_H4–J_H rearrangement on the second allele [11]. In addition, NALM1 cells can be induced to rearrange *IGK* and *IGL* light chain V- and J-gene segments upon inhibition of BCR-ABL1 kinase activity by STI571 [12]. Finally, NALM1 cells express the pan-B cell marker CD19 [11–13] but myeloid lineage markers (including CD13, CD14, GM-CSFR and IL3R α) are absent or expressed only at low levels. By genotypic and phenotypic definition, these data collectively identify NALM1 cells as a B lymphoid cell line. Likewise, BV173 cells carry rearranged *IGH* V region genes and express CD19, in apparent contradiction to their widespread designation as “CML cell line” [16–29].

Bonaventure Ndikung Bejeng Soh
 Florian Klein
 Niklas Feldhahn
 Markus Mischen

Laboratory für Molecular Stem Cell Biology
Heinrich-Heine-Universität Düsseldorf
Gebäude 14.80, Moorenstraße, 5
40225 Düsseldorf
Germany

References

- [1] Calabretta B, Perrotti D. The biology of CML blast crisis. *Blood* 2004;103:4010–22.
- [2] Niki M, Di Cristofano A, Zhao M, Honda H, Hirai H, Van Aelst L, Cordon-Cardo C, Pandolfi PP. Role of dok-1 and dok-2 in leukemia suppression. *J Exp Med* 2004;200:1689–95.
- [3] Yasuda T, Shirakata M, Iwama A, Ishii A, Ebihara Y, Osawa M, Honda K, Shinohara H, Sudo K, Tsuji K, Nakauchi H, Iwakura Y, Hirai H, Oda H, Yamamoto T, Yamanashi Y. Role of dok-1 and dok-2 in myeloid homeostasis and suppression of leukemia. *J Exp Med* 2004;200:1681–7.
- [4] Jamieson CH, Ailles LE, Dylla SJ, Muijtens M, Jones C, Zehnder JL, Gotlib J, Li K, Manz MG, Keating A, Sawyers CL, Weissman IL. Granulocyte-macrophage progenitors as candidate leukemic stem cells in blast-crisis CML. *N Engl J Med* 2004;351:657–67.
- [5] Laurent E, Talpaz M, Kantarjian H, Kurzrock R. The BCR gene and Philadelphia chromosome-positive leukemogenesis. *Cancer Res* 2001;61:2343–55.
- [6] Huettner CS, Zhang P, Van Etten RA, Tenen DG. Reversibility of acute B-cell leukaemia induced by BCR-ABL1. *Nat Genet* 2000;24:57–60.
- [7] Lugo TG, Pendergast AM, Muller AJ, Witte ON. Tyrosine kinase activity and transformation potency of bcr-abl oncogene products. *Science* 1990;247:1079–82.
- [8] Hu Y, Liu Y, Pelletier S, Buchdunger E, Warmuth M, Fabbro D, Hallek M, Van Etten RA, Li S. Requirement of Src kinases Lyn, Hck and Fgr for BCR-ABL1-induced B-lymphoblastic leukemia but not chronic myeloid leukemia. *Nat Genet* 2004;36:453–61.
- [9] Sill H, Goldman JM, Cross NC. Homozygous deletions of the p16 tumor-suppressor gene are associated with lymphoid transformation of chronic myeloid leukemia. *Blood* 1995;85:2013–6.
- [10] Drexler HG, MacLeod RA, Uphoff CC. Leukemia cell lines: in vitro models for the study of Philadelphia chromosome-positive leukemia. *Leuk Res* 1999;23:207–15.
- [11] Klein F, Feldhahn N, Harder L, Wang H, Wartenberg M, Hofmann WK, Wernet P, Siebert R, Muschen M. The BCR-ABL1 kinase bypasses selection for the expression of a pre-B cell receptor in pre-B acute lymphoblastic leukemia cells. *J Exp Med* 2004;199:673–85.
- [12] Klein F, Feldhahn N, Mooster JL, Sprangers M, Hofmann WK, Wernet P, Wartenberg M, Müschen M. Tracing the pre-B to immature B cell transition in human leukemia cells reveals a coordinated sequence of primary and secondary IGK gene rearrangement, IGK deletion, and IGL gene rearrangement. *J Immunol* 2005;174:367–75.
- [13] Feldhahn N, Klein F, Mooster JL, Hadweh P, Sprangers M, Wartenberg M, Bekhite MM, Hofmann WK, Herzog S, Jumaa H, Rowley J, Müschen M. Mimicry of a constitutively active pre-B cell receptor: Aberrant splicing links BTK to BCR-ABL1 in pre-B lymphoblastic leukemia. *J Exp Med* 201 (in press).
- [14] Sonta SI, Minowada J, Tsubota T, Sandberg AA. Cytogenetic study of a new Ph1-positive cell line (NALM-1). *J Natl Cancer Inst* 1977;59:833–7.
- [15] Drew SI, Carter BM, Terasaki PI, Naiem F, Nathanson DS, Abromowitz B, Gale RP. Cell surface antigens detected on mature and leukemic granulocytic populations by cytotoxicity testing. *Tissue Antigens* 1978;12:75–86.
- [16] Westbrook CA, Rubin CM, Carrino JJ, Le Beau MM, Bernards A, Rowley JD. Long-range mapping of the Philadelphia chromosome by pulsed-field gel electrophoresis. *Blood* 1988;71:697–702.
- [17] Van Denderen J, Hermans A, Meeuwssen T, Troelstra C, Zegers N, Boersma W, Grosveld G, van Ewijk W. Antibody recognition of the tumor-specific bcr-abl joining region in chronic myeloid leukemia. *J Exp Med* 1989;169:87–98.
- [18] Posner MR, Santos DJ, Elboim HS, Tumber MB, Frackelton AR Jr. Human monoclonal antibodies reactive with human myelomonocytic leukemia cells. *Cancer Res* 1989;49:1665–70.
- [19] Zhang W, Hittelman W, Van N, Andreeff M, Deisseroth A. The phosphorylation of retinoblastoma gene product in human myeloid leukemia cells during the cell cycle. *Biochem Biophys Res Commun* 1992;184:212–6.
- [20] Bi S, Hughes T, Bungey J, Chase A, de Fabritiis P, Goldman JM. p53 in chronic myeloid leukemia cell lines. *Leukemia* 1992;6:839–42.
- [21] Sattler M, Verma S, Byrne CH, Shrikhande G, Winkler T, Algate PA, Rohrschneider LR, Griffin JD. BCR/ABL directly inhibits expression of SHIP, an SH2-containing polyinositol-5-phosphatase involved in the regulation of hematopoiesis. *Mol Cell Biol* 1999;19:7473–80.
- [22] Vieira SA, Deininger MW, Sorour A, Sinclair P, Foroni L, Goldman JM, Melo JV. Transcription factor BACH2 is transcriptionally regulated by the BCR/ABL oncogene. *Genes Chromosomes Cancer* 2001;32:353–63.
- [23] Fajkusova L, Fajkus J, Polackova K, Fulnecek J, Dvorakova D, Krahulcova E. Detailed mapping of methylcytosine positions at the CpG island surrounding the Pa promoter at the bcr-abl locus in CML patients and in two cell lines, K562 and BV173. *Blood Cells Mol Dis* 2000;26:193–204.
- [24] Dazzi F, Capelli D, Hasserjian R, Cotter F, Corbo M, Poletti A, Chinswangwatanakul W, Goldman JM, Gordon MY. The kinetics and extent of engraftment of chronic myelogenous leukemia cells in non-obese diabetic/severe combined immunodeficiency mice reflect the phase of the donor's disease: an in vivo model of chronic myelogenous leukemia biology. *Blood* 1998;92:1390–6.
- [25] Rizzo MG, Zepparoni A, Cristofanelli B, Scardigli R, Crescenzi M, Blandino G, Giuliani S, Ferrari S, Soddu S, Sacchi A. Wt-p53 action in human leukaemia cell lines corresponding to different stages of differentiation. *Br J Cancer* 1998;77:1429–38.
- [26] Smetsers TF, Linders EH, van de Locht LT, de Witte TM, Mensink EJ. An antisense Bcr-Abl phosphodiester-tailed methylphosphonate oligonucleotide reduces the growth of chronic myeloid leukaemia patient cells by a non-antisense mechanism. *Br J Haematol* 1997;96:377–81.
- [27] Scheid S, Heinzinger M, Waller CF, Lange W. Bcl-2 mRNA-targeted ribozymes: effects on programmed cell death in chronic myelogenous leukemia cell lines. *Ann Hematol* 1998;76:117–25.
- [28] Maekawa T, Kimura S, Hirakawa K, Murakami A, Zon G, Abe T. Sequence specificity on the growth suppression and induction of apoptosis of chronic myeloid leukemia cells by BCR-ABL anti-sense oligodeoxynucleoside phosphorothioates. *Int J Cancer* 1995;62:63–9.
- [29] O'Brien SG, Kirkland MA, Melo JV, Rao MH, Davidson RJ, McDonald C, Goldman JM. Antisense BCR-ABL oligomers cause non-specific inhibition of chronic myeloid leukemia cell lines. *Leukemia* 1994;8:2156–62.
- [30] Pelz AF, Weillepp G, Wieacker PF. Re-analysis of the cell line NALM-1 karyotype by GTG-banding, spectral karyotyping, and whole chromosome painting. *Cancer Genet Cytogenet* 2005;156:59–61.